



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Exposure to diesel exhaust induces changes in EEG in human volunteers

Citation for published version:

Cruts, B, van Etten, L, Toernqvist, H, Blomberg, A, Sandstrom, T, Mills, NL & Borm, PJA 2008, 'Exposure to diesel exhaust induces changes in EEG in human volunteers', *Particle and Fibre Toxicology*, vol. 5, 4, pp. -. <https://doi.org/10.1186/1743-8977-5-4>

Digital Object Identifier (DOI):

[10.1186/1743-8977-5-4](https://doi.org/10.1186/1743-8977-5-4)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Particle and Fibre Toxicology

Publisher Rights Statement:

© 2008 Crüts et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Short report

Open Access

Exposure to diesel exhaust induces changes in EEG in human volunteers

Björn Crüts¹, Ludo van Etten¹, Håkan Törnqvist², Anders Blomberg², Thomas Sandström², Nicholas L Mills³ and Paul JA Borm^{*1}

Address: ¹Centre of Expertise in Life Sciences, Zuyd University, Heerlen, The Netherlands, ²Department of Respiratory Medicine and Allergy, University of Umeå, Sweden and ³Centre for Cardiovascular Sciences, The University of Edinburgh, UK

Email: Björn Crüts - b.cruts@home.nl; Ludo van Etten - l.van.etten@hszuyd.nl; Håkan Törnqvist - hakan.tornqvist@lung.umu.se; Anders Blomberg - anders.blomberg@lung.umu.se; Thomas Sandström - thomas.sandstrom@lung.umu.se; Nicholas L Mills - nick.mills@ed.ac.uk; Paul JA Borm^{*} - p.borm@hszuyd.nl

^{*} Corresponding author

Published: 11 March 2008

Received: 14 November 2007

Particle and Fibre Toxicology 2008, 5:4 doi:10.1186/1743-8977-5-4

Accepted: 11 March 2008

This article is available from: <http://www.particleandfibretoxicology.com/content/5/1/4>

© 2008 Crüts et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Ambient particulate matter and nanoparticles have been shown to translocate to the brain, and potentially influence the central nervous system. No data are available whether this may lead to functional changes in the brain.

Methods: We exposed 10 human volunteers to dilute diesel exhaust (DE, 300 µg/m³) as a model for ambient PM exposure and filtered air for one hour using a double blind randomized crossover design. Brain activity was monitored during and for one hour following each exposure using quantitative electroencephalography (QEEG) at 8 different sites on the scalp. The frequency spectrum of the EEG signals was used to calculate the median power frequency (MPF) and specific frequency bands of the QEEG.

Results: Our data demonstrate a significant increase in MPF in response to DE in the frontal cortex within 30 min into exposure. The increase in MPF is primarily caused by an increase in fast wave activity (β2) and continues to rise during the 1 hour post-exposure interval.

Conclusion: This study is the first to show a functional effect of DE exposure in the human brain, indicating a general cortical stress response. Further studies are required to determine whether this effect is mediated by the nanoparticles in DE and to define the precise pathways involved.

Introduction

Several epidemiological studies have identified diesel exhaust as an important component in determining the adverse health effects of particulate matter (PM) air pollution [1,2]. The molecular toxicity of diesel exhaust [3], is suggested to include oxidative stress-mediated inflammation, through particle surface, polycyclic aromatic hydrocarbons and redox active metals. Inflammation is

considered to be central to both the pulmonary and systemic adverse health effects of diesel through environmental PM exposure [4]. Over the past decades several, several studies have suggested that inhaled nanoparticles are able to translocate to the brain via the olfactory nerves [5,6], where they have been associated with inflammatory changes at sites of deposition [7,8]. Passage to the brain is of particular concern since nanoparticles are potent induc-

ers of oxidative stress [4,9] and the brain is very sensitive to damage caused by oxidative stress [10]. Oxidative stress has been implicated in the pathogenesis of neurodegenerative diseases such as Parkinson's and Alzheimer's disease and it is conceivable that the long-term effects of PM exposure might include a decrease in cognitive function [11]. Exposure to PM in an experimental mouse model resulted in widespread activation of pro-inflammatory cytokines in the brain [7].

First epidemiological evidence for a functional effect of PM on brain function was presented in recent study that suggests an association between black carbon levels in the environment and cognitive development of children [12]. As a first experimental step to investigate functional effects of PM and its ultrafine fraction, we studied the short-term changes in brain activity induced by exposure to diesel exhaust. We exposed human volunteers for one hour to diesel exhaust (DE, 300 µg/m³) and filtered air and followed their brain activity using quantitative electroencephalography (QEEG) at different sites of the scalp during and after one hour after exposure. The exposure protocol reflects a peak exposure that may occur in environmental and occupational exposure to diesel exhaust.

Methods

Ten subjects (all male, mean age: 26 years, range: 18–39 years; free of neurological or psychopathological impairments) were exposed to diesel exhaust and filtered air during one hour (sham condition) in a blinded randomized crossover design, separated by a period of two to four days. Prior to the start of each exposure, resting brain activity was measured using QEEG outside the exposure chamber during 3-minute eyes open (Table 1) and 3-minute eyes closed periods. After the 1 hour exposure, EEG measurements were continued for another hour. Participants gave written informed consent and the study was approved by the Ethics Committee of the University Hospital of Umeå. Diesel exhaust was produced by a Volvo Diesel engine (Volvo TD45, 4.5 L, 4 cylinders, 680 rpm) as described previously [13], leading to 1.2×10^6 suspended particles/cm³ (300 µg/m³) and gaseous pollutant levels of nitrogen dioxide (NO₂, 1.6 ppm, Nitrogen oxide (NO), 4.5 ppm, carbon monoxide (CO), 7.5 ppm and total hydrocarbons of 4.3 ppm.

During the EEG measurement, subjects rested in silence and performed no exercise to avoid interference on the QEEG measurement. Subjects sat in an upright position during the exposures, and were allowed to read a book to avoid boredom. EEG was continuously recorded from 8 electrode sites on the scalp according to the international 10–20 system: frontal pole (Fp1, Fp2), frontal (F3, F4), central (C3, C4) and parietal (P3, P4) at a sample rate of 500 Hz. The frequency content of the filtered EEG signals

Table 1: Individual absolute values of the median power frequency (MPF) at frontal locations (Fp1) and central location (C3) at pre- and post-exposure measurements (3 min each).

Nr	MPF at Fp1 (Hz)				MPF at C3 (Hz)			
	Sham		Diesel		Sham		Diesel	
	pre	post	pre	post	pre	post	pre	post
1	5.7	7.6	6.4	8.0	6.5	8.7	6.9	9.4
2	7.1	8.3	8.7	12.6	7.8	6.1	9.5	11.2
3	7.8	8.5	7.4	8.4	9.7	8.4	6.7	10.2
4	8.3	7.7	8.7	9	11.5	9.9	9	9.6
5	9.2	9.1	9.8	8.4	10.5	12.3	9.7	9.2
6	7.1	10.7	8.9	11	11	11	9.0	11.9
7	15.3	12.7	9.3	14.4	15.1	13.6	7.9	9.1
8	10.2	11	7.4	11.3	7.6	9.1	8.3	9.3
9	7.5	7.3	6.6	7.1	7	6.3	7.2	8.0
10	6.7	7	8.9	11.2	11.2	12.3	7.4	9.1
Mean	8.5	9.0	8.2	10.1^a	9.9	9.8	8.2	9.7^a
SD	2.7	1.9	1.2	2.3	2.6	2.6	1.1	1.1

Measurements were done before entering the exposure chamber and after completion of post-exposure interval, in exposure conditions sham and diesel.

a) Significantly different from pre-exposure, $P < 0.02$, Wilcoxon paired test

was calculated using Fast Fourier Transform (FFT) of 15 seconds intervals. From these intervals the Median Power Frequency (MPF) and spectral bands of the EEG were calculated. A second more detailed analysis divided the power spectrum into distinct frequency bands, including delta (1–3.5 Hz), theta (3.5–7.5 Hz), alpha (7.5–12 Hz), beta1 (12–20 Hz) and beta2 (20–32 Hz) bands. Changes in MPF and spectral bands during and after exposure are first presented as time series. From these time series, the mean MPF and spectral bands per hour and amplitudes of the first and last 5 minutes per exposure or post-exposure interval were calculated. Significance levels of differences ($P < 0.05$) between the first and last 5 minutes were compared between diesel and sham conditions by applying paired Wilcoxon tests to group results.

Results

Changes in Median Power Frequency (MPF)

Baseline measurements of MPF in the exposure chamber did not differ from pre-exposure measurements obtained outside the exposure chamber (data not shown). This shows that there is no acute effect of being in the exposure chamber, such as the presence of a pungent smell remaining after diesel sessions. There were also no significant differences at any time in MPF between subjects during the first 5 minutes of sham or diesel exposure. From 30 minutes into exposure we observed a slow increase in MPF of all subjects within the exposure chamber which was stronger in diesel exhaust exposure as compared to filtered air (Fig. 1). The effect on MPF was first observed and most




Figure 1

Time series of MPF at the left frontal cortex (F3).

Changes in MPF are represented as a function of time in a superposed epoch analysis graph for all subjects combined. Values for MPF are visualized for both diesel (blue line) and sham condition (black line) during the exposure and post exposure period. Both lines represent 5 minute moving averages of the original signal. MPF values during the diesel exposure increase and remain elevated during the post exposure period condition.

pronounced at the frontal electrode sites Fp1, Fp2, F3, and F4 (Fig. 2) and the average MPF at Fp1 and Fp2 was significantly different at the end of the diesel exhaust exposure compared to the start of exposure and to the sham conditions ($P < 0.05$ for both, Table 1). Interestingly the effect further increased during the diesel post-exposure period, when subjects were removed from the exposure chamber. This led to a significant difference between the 3 minute pre- and post-exposure measurements of MPF at the start and end of the post-exposure period both frontal (Fp1) and at the central (C3) location (Table 1). In addition the change over the exposure intervals, calculated from the difference between the first and last 5 minutes into the chamber, was significantly different between sham and diesel exposure (Table 2). This again indicated that tasks or factors present in the exposure chamber are unlikely to play a role in this effect on EEG. In addition, during post-exposure the effect on MPF spread in the other frontal F3 and F4 electrode sites leading to significant differences between diesel and sham (Table 2). Although slower changes in MPF were observed at the central (C3, C4) and parietal (P3, P4) electrode sites during the diesel (Fig 2), at the end of exposure the mean MPF at post-exposure measurement was also significantly different from pre-exposure (Table 1).

Analysis of EEG frequency bands

Subsequently we extended the data-analysis to determine the effects of diesel exhaust on specific frequency bands

from the EEG signal. The increased MPF at frontal electrode sites (Fp1, Fp2, F3, F4), is largely explained by an effect on fast wave activity (20–32 Hz, also denominated as β_2 activity). During diesel exposure, β_2 activity increased significantly compared to sham exposure. A marked increase in the β_2 signal was observed at the end of the diesel exposure interval in 7 out of 10 subjects (Fig. 3). The resulting change remained throughout the entire post exposure period. However, no such change in β_2 activity was observed at sham exposure (Fig. 3). Similar changes in fast wave activity were not observed at central and parietal electrode sites. Time variations of all frequency bands were analyzed using short-term Fourier-transformation, which revealed large variations in fast wave activity, with the largest variations over time in β_2 values (data not shown). β_1 -activity (15–20 Hz) also increased during diesel exposure at the frontal cortex, but this rise did not reach statistical significance (data not shown). Delta, theta and alpha activity showed no significant alterations during the diesel and post exposure periods compared to sham conditions (Fig. 3).

Discussion

This is the first study to demonstrate functional changes in brain activity as a result of exposure to diesel exhaust (DE) in human subjects. Our data demonstrate a delayed response to DE in the frontal cortex, characterized by an increase of median power frequency (MPF) and fast wave activity (21–32 Hz) in the EEG. These findings suggest an increased activity of the left frontal cortex during and after DE, but may be mediated by a number of different pathways.

As the first study of its kind interpretation of these findings are limited by a number of factors. Perhaps the most important limitation of our study is the fact that exposure to diesel exhaust is a mixture of combustion derived nanoparticles (CDNP) and exhaust gases (hydrocarbons, CO, NO_x). A direct effect of nanoparticles after or upon the translocation of the nanoparticles in DE may play a role, but no direct evidence is presented for this explanation. It is known that upon inhalation a large fraction of nanoparticles will deposit in the nasal cavity [14]. From there DEP, but also carrier constituents such as polycyclic aromatic hydrocarbons (PAHs) and redox metals in DE may migrate through the epithelium to the olfactory bulb [5,6,8] and cause mild inflammation and toxicity [7,8]. Uptake of MnO₂ nanoparticles in the olfactory bulb of rats has been reported beyond 6 hours after exposure, although no earlier time or inflammatory markers were investigated [7,8]. Therefore we think it is unlikely that the effect is induced by a toxic or inflammatory effect of diesel particles in the brain. In addition, gases such as CO, NO_x and hydrocarbons may have mediated changes in EEG through vagal reflexes in the airways. Endogenous




Figure 2
MPF values per electrode localisation. The figures represent the 8 electrode localizations, from the frontal areas at the top (Fp1, Fp2) to the parietal areas at the bottom (P3, P4). Amplitudes of MPF are indicated by colors, ranging from deep blue (7 Hz) to red (12 Hz). MPF amplitudes are presented as mean values of the first and last 5 minutes of the exposure hour and post exposure hour. The highest values for MPF during diesel exposure are observed at the frontal polar electrode sites (Fp1, Fp2). At these sites MPF increases significantly during diesel exposure compared to sham exposure. Following the diesel exposure MPF continued to increase resulting in significant differences at the frontal polar and the frontal sites (Fp1, Fp2, F3, F4) compared to the post-sham exposure period.

carbon monoxide (CO) generated by the heme oxygenase system has been shown in numerous studies to play a role in cardiac and neurophysiologic responses, but little information is available whether these findings are also relevant in traffic or diesel exhaust exposure. Interestingly, diesel particles may activate a pro-inflammatory vagovagal reflex in rats which is reduced by atropine [15]. This suggests that diesel particles themselves stimulate vagal reflexes in the airways and may as such cause feedback to the brain. On the other hand no difference in heart rate

Table 2: Mean absolute change in median power frequency (MPF) at different locations during different exposure conditions.

Condition	Electrode locations							
	Fp1	Fp2	F3	F4	C3	C4	P3	P4
ΔDiesel	0,41*	0,34*	0,40	0,54	0,12	0,18	0,21	0,24
ΔSham	0,10	0,12	0,34	0,50	0,23	0,13	0,17	0,25
ΔDiesel post	0,46*	0,43*	0,41*	0,60*	0,23	0,20	0,23	0,30
ΔSham post	0,21	0,24	0,09	-0,20	0,18	0,15	0,21	0,19

Mean changes in the median power frequency between the first and last 5 minutes of all conditions are shown per electrode localization. * Significant differences between diesel and sham exposures, Wilcoxon paired test (P < 0.05).




Figure 3
Absolute beta2 amplitudes (A) and theta amplitudes (B) at the left frontal cortex (F3). Changes are represented as a function of time in a superposed epoch analysis graph for all subjects combined. Absolute power per frequency band is visualized for diesel (blue line) and sham condition (black line), combining the one hour exposure condition and subsequent post condition in one graph. Both lines represent a 5 minute moving average of the original signal. Beta2 (A) but not theta-amplitudes (B) are elevated during diesel exposure compared to sham exposure.

was seen between exposures, which would have occurred of the current diesel exhaust had induced a vagal reflex.

A further complication is that the exposure room was contaminated with a pungent smell and it cannot be excluded that the smell of diesel exhaust plays a role in the increased cortical arousal in the exposed subjects. However a confounding effect on our findings by the smell is unlikely for several reasons. First, subjects reported only discomfort due to smell in the first minutes while changes in EEG were only seen at the end of the exposure interval. Secondly, the effects on MPF continued to rise after subjects left the exposure chamber. Thirdly, the diesel odor was also present during the sham condition, because of alternating use of the exposure chamber for diesel and filtered air exposure. Finally, olfactory sensitization to chemical stimulation is usually noted as an increase of slow-wave activity within a few minutes of onset of expo-

sure [16]. This slow wave activity was unaffected during the entire study.

Interpretation of our findings is hampered by the lack of similar studies on such a time-scale. Usually EEG studies are done on short-term exposure or tasks, that last only a few minutes. Therefore we have to consider that the increased activity of the frontal cortex could also relate to lowered vigilance of the subjects, caused by weariness or getting bored during the exposure sessions. Indeed, increased β_2 values have been observed in combination with lowered vigilance during cognitive tasks [17] and are often regarded as compensation for lowered vigilance. However this is usually associated with increased levels of EEG slow wave activity [18], which was not observed in this study. In addition, no significant effect of β_2 fast-wave activity was seen in the same subjects during sham exposure. Therefore we conclude that decreased vigilance is unlikely to explain the observed effects. Other studies using EEG have linked frontal brain activity to various physiological and psychological findings. Increased β_2 levels have been reported in patients with neurological and psycho-pathological disorders, such as headache, post-traumatic stress disorder, burnout and traumatic brain injury, and are often regarded as indicators of cortical stress [19,20]. Although tempting, linking these effects of short-term functional changes in the frontal cortex to decreased cognitive function across assessments of verbal and nonverbal intelligence and memory constructs in epidemiological work [12] is simply not possible at this stage.

Ambient PM exposure has been associated with inflammation in the lung, exaggerated airways responses and increased morbidity as well as mortality in respiratory and cardiovascular diseases [1,2]. Exposure to diesel exhaust has been associated with similar effects, and has been used as a surrogate exposure in many experimental studies [3,13,15]. The observed effects of diesel exhaust on the EEG add up to the evidence that air pollution may exert its effects by a variety of pathways and may induce effects in the brain. Based on the onset, location, frequency profile and persistence of the response in EEG, we suggest that our findings are due to an effect of nanoparticles that slowly penetrate the brain or affect neurophysiologic signaling. However we can only speculate what these effects on the frontal cortex may mean for chronic exposure to PM and/or nanoparticles. Further studies are necessary to establish whether other nanoparticles induce this effect, and to explore the dose-response relationship and clinical implications of these novel findings. It needs to be realized that further human studies are confronted with serious limitations, since neurochemical work and/or taking samples of spinal fluid are not possible. Subjecting the subjects to non-invasive techniques for imaging (Mag-

netic Resonance Imaging, MRI) or measuring (Near Infra-red Resonance Spectroscopy, NIRS) blood flow and tissue damage are the options that are currently being explored to follow-up these human studies. Current work concentrates on the similar effects of artificially generated nanoparticles that are not contaminated with gases and other components.

Authors' contributions

BC and LE carried out the pilot-measurements and data processing of the EEG signals, and prepared and supervised the project on-site; HT, was responsible for the exposure assessment, AB and NM were the responsible medical doctors for the survey; TS and PB were the coordinators and planners of the study. The manuscript was written by BC and PB, but all authors read, corrected and approved the manuscript.

Acknowledgements

The authors would like to acknowledge Maastricht Instruments (Maastricht, NL) for providing the MPAQ system; Biometrisch Centrum (Gulpen, NL) for the EEG equipment and software, Erik Hinssen and Felipe Ernst (Zuyd University students) for their measurements of QEEG, Annika Johansson and Helena Bogseth (research nurses at University Hospital, Umeå, Sweden), and finally SKO, the British Heart Foundation (RG/05/003), The Swedish Lung-Heart Foundation and the Swedish National Air Pollution programme for supporting this research effort.

References

- Pope CA 3rd, Burnett RT, Thurston GD, Thun MJ, Calle EE, Krewski D, Godleski JJ: **Cardiovascular mortality and long-term exposure to particulate air pollution: epidemiological evidence of general pathophysiological pathways of disease.** *Circulation* 2004, **109**(1):71-77.
- Laden F, Neas LM, Dockery DW, Schwartz J: **Association of fine particulate matter from different sources with daily mortality in six U.S. cities.** *Environ Health Perspect* 2000, **108**:941-947.
- Donaldson K, Tran L, Jimenez LA, Duffin R, Newby DE, Mills N, MacNee W, Stone V: **Combustion-derived nanoparticles: a review of their toxicology following inhalation exposure.** *Part Fibre Toxicol* 2005, **2**:10.
- Donaldson K, Stone V, Borm PJ, Jimenez LA, Gilmour PS, Schins RP, Knaapen AM, Rahman I, Faux SP, Brown DM, MacNee W: **Oxidative stress and calcium signaling in the adverse effects of environmental particles (PM10).** *Free Radic Biol Med* 2003, **34**:1369-1382.
- Oberdörster G, Sharp Z, Atudorei V, Elder A, Gelein R, Kreyling W, Cox C: **Translocation of inhaled ultrafine particles to the brain.** *Inhal Toxicol* 2004, **16**:437-445.
- DeLorenzo A: **The olfactory neuron and the blood-brain barrier.** In *Taste and smell in vertebrates* Edited by: Wolstenholme G, Knight J. J. & A. Churchill, London; 1970:151-176.
- Campbell A, Oldham M, Becaria A, Bondy SC, Meacher D, Sioutas C, Misra C, Mendez LB, Kleinman M: **Particulate matter in polluted air may increase biomarkers of inflammation in mouse brain.** *Neurotoxicology* 2005, **26**:133-140.
- Elder A, Gelein R, Silva V, Feikert T, Opanashuk L, Carter J, Potter R, Maynard A, Ito Y, Finkelstein J, Oberdörster G: **Translocation of inhaled ultrafine manganese oxide particles to the central nervous system.** *Environ Health Perspect* 2006, **114**:1172-1178.
- Nel A, Xia T, Madler L, Li N: **Toxic potential of materials at the nanolevel.** *Science* 2006, **311**(5761):622-627.
- Halliwel B: **Oxidative stress and neurodegeneration: where are we now?** *J Neurochem* 2006, **97**:1634-1658.
- Calderon-Garciduenas L, Reed W, Maronpot RR: **Brain inflammation and Alzheimer's-like pathology in individuals exposed to severe air pollution.** *Toxicol Pathol* 2004, **32**(6):650-658.
- Suglia SF, Gryparis A, Wright RO, Schwartz J, Wright RJ: **Association of black carbon with cognition among children in a prospective birth cohort study.** *Am J Epidemiol* 2008, **167**(3):280-6.
- Mills NL, Tornqvist H, Gonzalez MC, Vink E, Robinson SD, Soderberg S, Boon NA, Donaldson K, Sandstrom T, Blomberg A, Newby DE: **Ischemic and Thrombotic Effects of Dilute Diesel-Exhaust Inhalation in Men with Coronary Heart Disease.** *New Engl J Med* 2007, **357**:1075-1182.
- ICRP: **Human Respiratory Model for Radiological Protection.** *Annals of The ICRP* 24, ICRP publication # 66.
- McQueen DS, Donaldson K, Bond J, McNeilly JD, Newman J, Barton NJ, Duffin R: **Bilateral vagotomy or atropine pre-treatment reduces experimental diesel-soot induced lung inflammation.** *Toxicol Appl Pharmacol* 2007, **219**(1):62-71.
- Bell IR, Szarek MJ, Dicenso DR, Baldwin CM, Schwartz GE, Bootzin RR: **Patterns of waking EEG spectral power in chemically intolerant individuals during repeated chemical exposures.** *Int J Neurosci* 1999, **97**(1-2):41-59.
- Smit AS, Eling PA, Coenen AM: **Mental effort affects vigilance enduringly: after-effects in EEG and behavior.** *Int J Psychophysiol* 2004, **53**(3):239-243.
- Coull JT: **Neural correlates of attention and arousal: insights from electrophysiology, functional neuroimaging and psychopharmacology.** *Prog Neurobiol* 1998, **55**(4):343-361.
- De Carlo L, Cavaliere B, Arnaldi C, Faggioli R, Soriani S, Scarpa P: **EEG evaluation in children and adolescents with chronic headaches.** *Eur J Pediatr* 1999, **158**:247-8.
- Thornton K: **The electrophysiological effects of a brain injury on auditory memory functioning. The QEEG correlates of impaired memory.** *Arch Clin Neuropsychol* 2003, **18**:363-378.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

